

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

EPA Reg. Number:

Date of Issuance:

9198-219

AUG 2 0 2013

Term of Issuance: Unconditional

Name of Pesticide Product: The Andersons 0.72% Granular Propiconazole

NOTICE OF PESTICIDE:

X Registration
X Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

The Andersons Lawn Fert. Div. Inc.

PO Box 119

Maumee, OH 43537

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted by email on August 15, 2013. EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. With this accepted labeling, all requirements set forth in the Reregistation Eligibility Decision for propiconazole have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Page 1 of 2

Signature of Approving Official:

Date:

AUG 2 0 2013

Shaja Joyner, Product Manager (20)

Fungicide Branch/Registration Division/OPP/OCSPP (7504P)

Notice of Pesticide Reregistration The Andersons 0.72% Granular Propiconazole EPA Reg. No. 9198-219 Page 2 of 2

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

If you have any questions or comments regarding this letter, please contact Marcel Howard at (703) 305-6784 or via e-mail at howard.marcel@epa.gov.

Enclosure:

Label stamped "Accepted"
PRB Label Review dated 04/25/2013
Acute Toxicity Review DP408736 dated 03/19/2013
Product Chemistry Review DP402732 dated 01/14/2013

The Andersons 0.72% Granular Propiconazole

Optional marketing information:





Dispersing Granule Technology



ACCEPTED

AUG 2 0 2013

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No.

9198-219

- Fungicide
- Broad spectrum and systemic disease control and suppression for turf
- Dispersing granule fungicide
- Easy to spread formula

Active Ingredient: 0.72% Propiconazole* Other Ingredients: 99.28% Total:

*CAS No. 60207-90-1

EPA Reg. No. 9198-219 EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Visit our website: www.andersonspro.com

Net Weight XX lbs.

The Andersons 0.72% Granular Propiconazole

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed, absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear goggles or face shield. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made from any waterproof material
- Shoes plus socks
- Goggles or face shield

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment, disposing of equipment wash waters or rinsate.

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Note to Physic	l lingested, induce emesis or lavage stomach. Treat symptomatically.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, shoes and socks, and chemical-resistant gloves made from any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow adults, children, or pets to enter the treated areas until dust has settled.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry secure place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide wastes may be acutely hazardous. Improper disposal is a violation of Federal law. Pesticide, mixtures, or equipment rinse water that cannot be chemically reprocessed must be disposed of according to applicable Federal, state or local procedures. Contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a systemic fungicide for use on turfgrasses for the control and suppression of:

Dollar spot	Sclerotinia homoeocarpa	Spring dead	Leptosphaeria korrae,
		spot	leptosphaeria narmari,
			Ophiosphaerelia herpotricha,
			Gaeumannomyces graminis
Brown patch	Rhizoctonia solani	Take-all patch	Gaeumannomyces graminis
Anthracnose	Colletotrichum graminicola	Leafspot	Biopolaris spp., Drechslera
	_	·	spp.
Red thread	Laetisaria fuciformis	Gray leafspot	Pyricularia grisea
Pink patch	Limonomyces roseipellis	Pink snowmold	Microdochium nivale
Rust	Puccinia graminis	Fusarium patch	Microdochium nivale
Powdery mildew	Erysiphe graminis	Gray snowmold	Typhula spp.
Stripe smut	Ustilago striiformis and	Yellow patch	Rhizoctonia cerealis
•	Urocystis agropyri	•	
Summer patch	Magnaporthe poae	Zoysia patch	Rhizoctonia solani
Necrotic ring	Leptosphaeria korrae		•
spot	•		

Do not apply this product through any type of irrigation system.

Use this product in a preventative disease control program. Properly calibrate the spreader to ensure thorough coverage. Apply after mowing. For control of foliar diseases, apply to wet/moist foliage. For control of soil-borne diseases, this product can be watered in after application. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval. For optimum turf quality and disease control, use this product in conjunction with turf management practices that promote good plant health and optimum disease control. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

Do not apply more than 20 lbs.product/1,000 sq. ft./calendar year.

Important: Bermudagrass can be sensitive to this product. Do not exceed 5 lbs./1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply this product to bermudagrass golf greens when temperatures exceed 90°F.

Note: Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.

Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Lbs. product per 1,000 sq. ft.	Lbs. product per Acre	Application Timing / Interval	Instructions
Dollar spot Sclerotinia homoeocarpa	2.5	109	14-28 days	Make no more than 3 consecutive applications for dollar spot before rotating to an alternative EPA-registered fungicide having a different mode of action.
Anthracnose Colletotrichum graminicola	2.5	109	14-28 days	Apply when conditions are favorable for disease development. When disease pressure is high, use shorter intervals.
Brown patch Rhizoctonia solani	2.5	109	14-21 days	Begin applications in May or June before disease is present. Under conditions of high temperatures and high humidity, use this product at shorter intervals.
Powdery mildew Erysiphe graminis Rust Puccinia graminis	2.5	109	14-28 days	Apply when conditions are favorable for disease development.
Red thread Laetisaria fuciformis Pink patch Limonomyces roseipellis	2.5	109	14-21 days	Apply when conditions are favorable for disease development.
Stripe smut Ustilago striiformis Urocystis agropyri	2.5	109	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray leafspot Pyricularia grisea	2.5	109	14 days	Apply when conditions are favorable for disease development.
Melting Out Leafspot Biopolaris spp., Drechslera spp.	2.5	109	14 days	Under light to moderate pressure, apply this product to reduce the severity of leaf spot and melting out caused by Helminthosporium-type pathogens.
Summer patch	2.5	109	14 days	Apply this product beginning in April. Use
Poa Patch Magnaporthe poae	5.	218	28 daÿs	the 5 lb:/1,000 sq. ft. rate on a 28 day schedule and the 2.5 lb. /1,000 sq. ft. rate on a 14 day schedule.
Take-all patch Gaeumannomyces graminis	2.5-5	109-218	Spring and Fall	Apply this product to reduce the severity of take-all patch. Make 1-2 fall applications in September and October or when night temperatures drop to 55°F and 1-2 spring applications in April and May, depending on local recommendations.
Spring dead spot Leptosphaeria korrae, leptosphaeria narmari, Ophiosphaerelia herpotricha, Gaeumannomyces graminis	5	218	30 days	Make 1-3 applications. For one application, apply in September or October. For multiple applications, begin applications in August.
Necrotic ring spot Leptosphaeria korrae	5	218	Fall or Spring	Apply in the fall and/or the early spring depending on local recommendations.

Disease	Lbs. product per 1,000 sq. ft.	Lbs. product per Acre	Application Timing / Interval	Instructions
Snowmold Gray Typhula spp. Pink Microdochium nivale	2.5-5	109-218	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. For optimum disease control, the 2.5 and 3.75 lb. rates should be used in conjunction with either PCNB or chlorothalonil at label rates.
Fusarium patch Microdochium nivale	2.5-5	109-218	Fall-Early Spring	Apply when conditions are favorable for disease development.
Yellow patch Rhizoctonia cerealis	3.75-5	163-218	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. If using a 3.75 lb. /1,000 sq. ft. rate, use in conjunction a registered contact fungicide at the label rate.
Zoysia patch, Large patch of zoysia Rhizoctonia solani	3.75-5	163-218	Early Fall	Make one application in the early fall (mid- September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing in your area.

Dichondra - Specific Diseases, Rates, and Application Timing

Diolionata opcomo Diocacco, itateo, ana Approation inimig				<u></u>
Disease	Lbs. product per 1,000 sq. ft.	Lbs. product per Acre	Application Timing / Interval	Instructions
Dichondra Rust Puccinia dichondrae	2.5	109	14-21 days	Apply when conditions are favorable for disease development.

Spreader Settings

These spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.

40 lbs. treats 16,000 sq. ft. at the 109 lbs./acre rate (2.50#/1,000 sq. ft.) 40 lbs. treats 10,600 sq. ft. at the 163 lbs./acre rate (3.75#/1,000 sq. ft.) 40 lbs. treats 8,000sq. ft. at the 218 lbs./acre rate (5.00#/1 000 sq. ft.)

SPREADER	GROUND OR PTO	WIDTH OF COVERAGE	RECOMMENDED RATE Per 1,000 sq. ft.			
SPREADER	SPEED		2.50# (0.78 lbs. ai/acre)	3.75# (1.17 lbs. ai/acre)	5.00# (1.57 lbs. ai/acre)	
Andersons AccuPro/SR- 2000 (cone X)	X mph	XX ft	×	x	x	
Andersons SS-R Drop	X mph	XX ft.	X	X	X	
Scotts R-8A (cone X)	X mph	XX ft	X	X.	X	
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm), Pattern III-A	X mph	XX.ft	×	×	х	
Vicon (03 Series)	X mph	XX ft	Χ .	X	X	

Note to EPA: actual spreader models may vary depending on market demand and availability

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. PO Box 119 Maumee, OH 43537 www.andersonspro.com

Date Printed	08/15/2013
Date Approved	2/5/08
Supersedes	8/1/06
Label Changes	RED label amendment
Filename	9198-219
Label Code	RVred08152013